



## Research report

# A microstructural analysis of schedule-induced polydipsia reveals incentive-induced hyperactivity in an animal model of ADHD<sup>☆</sup>



Javier Íbias<sup>a</sup>, Ricardo Pellón<sup>a</sup>, Federico Sanabria<sup>b,\*</sup>

<sup>a</sup> Laboratorios de Conducta Animal, Departamento de Psicología Básica I, Facultad de Psicología, Universidad Nacional de Educación a Distancia (UNED), C/ Juan del Rosal 10, Ciudad Universitaria, 28040 Madrid, Spain

<sup>b</sup> Department of Psychology, Arizona State University, P.O. Box 871104, Tempe, AZ 85287-1104, USA

## HIGHLIGHTS

- Similar rate of licking induced by FT 30-s schedule in SHR, Wistar, and WKY rats.
- FT 90-s lengthened drinking episodes in SHR, reduced rate of licking in Wistar, WKY.
- Schedule-induced drinking occurred in bouts separated by pauses.
- SHR episodes comprised shorter, more frequent bouts of drinking.

## ARTICLE INFO

### Article history:

Received 1 September 2014

Received in revised form 13 October 2014

Accepted 17 October 2014

Available online 27 October 2014

### Keywords:

Schedule-induced polydipsia  
Spontaneously hypertensive rat  
Hyperactivity  
Licking Bouts  
BERM

## ABSTRACT

Recent research has suggested that frequent short bursts of activity characterize hyperactivity associated with attention deficit hyperactivity disorder (ADHD). This study determined whether such pattern is also visible in schedule-induced polydipsia (SIP) in the spontaneously hypertensive rat (SHR), an animal model of ADHD. Male SHR, Wistar Kyoto (WKY) and Wistar rats were exposed to 40 sessions of SIP using a multiple fixed-time (FT) schedule of food delivery with FT 30-s and FT 90-s components. Stable performance was analyzed to determine the extent to which SIP-associated drinking is organized in bouts. The Bi-Exponential Refractory Model (BERM) of free-operant performance was applied to schedule-induced licks. A model comparison analysis supported BERM as a description of SIP episodes: licks were not produced at a constant rate but organized into bouts within drinking episodes. FT 30-s induced similar overall licking rates, latencies to first licks and episode durations across strains; FT 90-s induced longer episode durations in SHRs and reduced licking rate in WKY and Wistar rats to nearly baseline levels. Across schedules, SHRs made more and shorter bouts when compared to the other strains. These results suggest an incentive-induced hyperactivity in SHR that has been observed in operant behaviour and in children with ADHD.

© 2014 Elsevier B.V. All rights reserved.

## 1. Introduction

Recent research suggests that children with attention deficit hyperactivity disorder (ADHD) engage in frequent but short bouts of operant behaviour [1]. The spontaneously hypertensive rat (SHR), an animal model of ADHD [2,3], shows a similar pattern

of operant performance [4]. It thus appears that this pattern of behaviour constitutes a key component of the behavioural phenotype of ADHD.

The present study was aimed at determining whether frequent but short bouts of responding were also observed in SHRs in a non-operant behavioural preparation, schedule-induced polydipsia (SIP). SIP consists of excessive drinking that occurs on a schedule in which food is intermittently delivered, usually observed in partially food-deprived rats [5]. SIP can be produced in a wide range of behavioural schedules, including conditions in which animals are exposed to intermittent delivery of food regardless of their behaviour, such as fixed time (FT) schedules [6,7].

The principal characteristic of SIP is its excessiveness, which distinguishes it from others behaviours performed throughout inter-food intervals. Thus, SIP serves as the prototype of so-called

<sup>☆</sup> Research supported by Spanish Government Grant PSI2011-29399 (Ricardo Pellón, Principal Investigator) (Ministerio de Economía y Competitividad, Secretaría de Estado de Investigación, Desarrollo e Innovación). Javier Íbias was under a UNED predoctoral research grant that supported 3 months stay at Arizona State University. Federico Sanabria was supported by Grant MH094562 (National Institutes of Health, United States).

\* Corresponding author. Tel.: +1 4809654687; fax: +1 4809658544.  
E-mail address: [federico.sanabria@asu.edu](mailto:federico.sanabria@asu.edu) (F. Sanabria).

adjunctive behaviour ([8]; also referred to as interim behaviour [9]). In relation to ADHD, SIP successfully selects among experimental subjects those that display characteristics associated with excess behaviour [10,11] and other disorders associated with deficient impulse control [12,13], such as substance abuse [14], obsessive compulsive behaviour [15–19], and schizophrenia [20,21].

Comparisons of SIP between SHR, Wistar and Wistar Kyoto (WKY) rats have shown that (a) there is no significant difference in SIP among strains when the inter-food interval is short (30 s), but (b) whereas SIP declines with longer inter-food intervals (approximately 60 s) in WKY and Wistar rats, SIP in SHR remains relatively high [22,23]. Reduced SIP in control strains may occur because of (a) a reduction in the proportion of inter-food intervals that contain drinking episodes, (b) an increase in the latency to initiate drinking episodes, (c) a reduction in the duration of drinking episodes, and/or (d) a reduction in the rate of licking during a drinking episode. To the extent that licking rate is, like operant behaviour, organized in bouts [24,25], licking rate may decline because (a) licking bouts occur less often, (b) licking bouts are shorter, and/or (c) licking rate within bouts declines.

The present study aimed at examining these potential differences in SIP performance across three strains of rats, SHR, Wistar and WKY, and two food schedules, FT 30-s and FT 90-s. The organization of drinking episodes in bouts of licks was of particular interest, not only because of its ostensible link to ADHD, but also because parameters of bout organization have been associated with motor, motivational and learning variables [25]. Therefore, differences in parameter estimates across strains may suggest novel hypotheses about the behavioural mechanisms that underlie the differences in performance across strains, as well as on the behavioural mechanisms involved in SIP and related phenomena.

## 2. Method

### 2.1. Subjects

Twenty-four male rats of three strains – 8 SHR, 8 WKY and 8 Wistar – obtained from Charles River Laboratories (Lyon, France) were used. On arrival at the laboratory, animals were 10 weeks old; they were housed in groups, in an environmentally controlled room with a 12-h light–dark cycle (light from 08:00 to 20:00 h), ambient temperature of 17–23 °C, and 60% relative humidity. Once habituated to the animal facility, rats were housed singly in 18 cm × 32.5 cm × 20.5 cm transparent Plexiglas cages, with a metal-grid detachable roof that allowed for food to be deposited and a water bottle to be fitted.

Rats were 12 weeks old at the start of the experiment. Their average weights were, for SHR, 292 g (range: 277–302 g); for WKY, 339 g (range: 306–361 g); for Wistar, 373 g (range: 359–384 g). Weights were reduced to 80–85% of free-feeding weight by a controlled diet, and then maintained throughout the experiment in proportion to standard growth curves for each strain. Each rat was weighed daily before the experimental session. Twenty minutes after the experimental session, each animal received the appropriate food supplement to maintain its weight within the criterion-based range.

### 2.2. Apparatus

Eight Letica LI-836 conditioning chambers, measuring 29 cm × 24.5 cm × 35.5 cm, enclosed in soundproofed housing, equipped with its own ventilation and a small observation window at the front. The front panel of each conditioning chamber was made of aluminium, the left wall of transparent Plexiglas and the remaining walls of black Plexiglas. On the exterior of the

chamber' right-hand wall, a water bottle was fitted, and the rat had access to the spout from the interior of the chamber, through a 3.2 cm × 3.9 cm aperture in the wall, situated 20 cm from the front panel and 7 cm from the floor. The spout was placed 2 cm towards the interior of the aperture to allow for licks rather than continuous drinking. Contact between the animal's tongue and the metal spout completed the electric circuit between the 12-bar metal grid that served as the floor and the water-bottle spout. Licks were recorded using a MED-PC-IV application under a Windows XP environment. Forty-five mg food pellets were dispensed (Bio-Serv, Frenchtown, NJ, USA) in an aperture in the chamber's front wall situated 3.7 cm from the floor, between the panel's two levers, which were retracted throughout the experiment. The chambers were lit by two 3 W lamps situated on the front panel at either side of the food hopper and by an indirect 25 W light fitted to the interior of the soundproof housing that insulated each chamber. Exterior noise was masked by a fan that produced an ambient noise of approximately 60 dB in each chamber. At the top of the front wall of each chamber was a speaker that produced sound signals when necessary.

### 2.3. Procedure

The SIP procedure was carried out using a multiple FT schedule, in which a food pellet was delivered at regular intervals regardless of the animal's behaviour. Every experimental session contained two schedule components, FT 30-s and FT 90-s, and the first component to start each session was determined randomly. Rats were given 20 food pellets/session in each component. One of the components was always signalled with a continuous tone of 60 dB (while the other was signalled by its absence); assignment of tone-signalled component was randomized across animals. The experimental chamber lights were turned off for 60 s between FT components.

Forty experimental sessions were conducted, until SIP performance was stable in both schedules in terms of lick rate (licks/min) and volume of water consumed. Data from the last 8 sessions was used for the estimation of bout parameters. All data were collected with a resolution of 20 ms.

### 2.4. Estimation of bout parameters


Bout-parameter estimation was conducted only on rats that, during the last 8 sessions, drank more than the average daily level of prandial water consumption (2.75 ml). A total of 19 rats (all SHR rats, 6 WKY rats and 5 Wistar rats) were included in the analysis.

The following measures were obtained from each trial in which a rat produced at least two licks: latency to the first lick, duration of the drinking episode (i.e., time between first and last lick), and 3 parameters of the distribution of inter-lick intervals (ILIs): the rate at which licking bouts were initiated ( $b$ ), the rate of licking within bouts ( $w$ ), and the probability that a lick was not the last one in a bout [ $p$ ; mean bout length =  $1/(1 - p)$ ]. These parameters were estimated using the Biexponential Refractory Model (BERM) of free-operant performance [26],

$$p(\text{ILI} = t | t < \delta) = 0$$

$$p(\text{ILI} = t | t \geq \delta) = pw \exp[-w(t - \delta)] + (1 - p)b \exp[-b(t - \delta)] \quad (1)$$

This equation indicates that there is a minimum time between consecutive licks,  $\delta$ , and that ILIs longer than  $\delta$  are sampled with probability  $p$  from an exponential distribution with mean  $(1/w) + \delta$  (the within-bout lick rate), and with probability  $1 - p$  from another exponential distribution with mean  $(1/b) + \delta$  (the bout-initiation rate). Estimates of  $p$ ,  $w$  and  $b$  were obtained using the method of least squares [27], with  $\delta$  fixed at 10 ms (half of the data resolution)



**Fig. 1.** Mean ( $\pm$ SEM) licks/min over acquisition sessions of SIP in FT 30-s (A) and FT 90-s (B). In FT 30-s, SHR produced significantly more licks/min than Wistar rats in sessions 4 through 11, and more than WKY in sessions 4 through 31 ( $p < 0.05$  for all comparisons). In FT 90-s, SHR produced significantly more licks/min than the other two strains from session 2 onwards ( $p < 0.05$ ).

for all rats. Goodness-of-fit of BERM parameters was computed (a) as the proportion of variance in the empirical survival function, expressed in log units [28], accounted for by the model (PVAf [29]), and (b) as the improvement in likelihood (probability of the model given the data) relative to a single shifted-exponential distribution, correcting for the 2 extra free parameters in Eq. (1) (the mixture weight  $p$  and the parameter of the second exponential). The latter computation was conducted using  $\Delta$ AICc [30].

### 2.5. Statistical analysis

Licks per minute during the first 40 sessions were analyzed using a  $3 \times 2 \times 40$  mixed-design ANOVA, with factors *strain* (SHR, Wistar and WKY), *schedule* (FT 30- and 90-s), and *session* (each of 40 sessions). The following dependent measures were analyzed in the last 8 experimental sessions: the proportion of trials (i.e., inter-food intervals) with a drinking episode (2 or more licks), the median latency to start drinking episodes, the median duration of drinking episodes, and the mean number of bouts emitted per episodes (estimated from BERM parameters). Analysis was conducted using a  $3 \times 2$  mixed-design ANOVA with factors *strain* and *schedule*. PVAf-weighted, log-transformed BERM parameter estimates were compared between groups using a similar ANOVA (the log transform follows suggestions that BERM parameters are log-normally distributed across rats; see [31]).

Post hoc comparisons were carried out using pairwise comparisons with a Bonferroni correction for  $p$ . The lowest  $p$ -value reported in every analysis was 0.01, with  $\alpha = .05$ . All analyses were conducted using SPSS 19 © Software.

## 3. Results

### 3.1. SIP acquisition

Fig. 1A shows mean ( $\pm$ SEM) licks/min throughout 40 sessions of SIP acquisition for the three strains of rats in the FT 30-s schedule. A significant *strain*  $\times$  *session* interaction effect was observed [ $F(78, 819) = 2.146, p < 0.01$ ]. Post hoc comparisons revealed that,

in sessions 4–11, SHRs produced significantly more licks/min than Wistar and WKY rats ( $p < 0.05$  for all comparisons). This difference persisted for SHR vs. WKY rats until session 31 ( $p < 0.05$  for all comparisons). No significant difference in licks/min among strains was observed in the last 9 of the first 40 sessions of FT 30-s training.


Fig. 1B shows mean ( $\pm$ SEM) lick/min throughout 40 sessions of SIP acquisition for the three strains of rats in the FT 90-s schedule. A significant *strain*  $\times$  *session* interaction effect was observed [ $F(78, 819) = 3.265, p < 0.01$ ]. Post hoc comparisons revealed that, from session 2 onwards, SHRs produced significantly more licks/min than Wistar and WKY rats ( $p < 0.05$  for both comparisons).

The analyses reported in the remainder of this section are confined to the last 8 training sessions. Mean  $\pm$ SD volumes of water consumed throughout these sessions were  $13.0 \pm 0.6$  ml for SHR,  $10.5 \pm 0.9$  ml for Wistar, and  $7.6 \pm 0.8$  ml for WKY. No significant differences in lick efficiency (volume of water per lick) were observed between strains. Mean  $\pm$ SD estimations of lick efficiency, in licks/ml, were:  $110.5 \pm 10.7$  for SHR,  $94.6 \pm 13.6$  for Wistar, and  $117.6 \pm 12.4$  for WKY.

Fig. 2A shows mean ( $\pm$ SEM) proportions of trials with at least 2 licks. An ANOVA of this dependent measure resulted in a significant *strain*  $\times$  *schedule* interaction effect [ $F(2, 16) = 8.317, p < 0.01$ ]. Post hoc comparisons revealed that SHR rats had significantly more drinking episodes than the other strains in both FT schedules ( $p < 0.01$  in all cases). These comparisons also showed a substantial decrease in the number of drinking episodes in Wistar and WKY rats in FT 90-s relative to FT 30-s schedules ( $p < 0.01$  for both comparisons).

Fig. 2B shows mean ( $\pm$ SEM) median latencies to first lick in all FT schedules for all groups. An ANOVA revealed a significant *strain*  $\times$  *schedule* interaction effect [ $F(2, 16) = 4.064, p < 0.04$ ]. Post hoc comparisons showed that latencies increased for all strains between FT 30-s and FT 90-s ( $p < 0.01$  for all comparisons).

Fig. 2C shows mean ( $\pm$ SEM) median durations of drinking episodes. An ANOVA of episode durations revealed a significant *strain*  $\times$  *schedule* interaction effect [ $F(2, 16) = 13.718, p < 0.01$ ]. Post hoc comparisons showed that SHR rats produced longer episode durations in FT 90-s than Wistar and WKY rats ( $p < 0.02$  and  $p < 0.04$ ,



**Fig. 2.** (A) Mean ( $\pm$ SEM) proportion of trials with at least 2 licks. (B) Mean ( $\pm$ SEM) median latency to first lick. (C) Mean ( $\pm$ SEM) median duration of drinking episodes (intervals between first and last lick). SHR produced more drinking episodes than Wistar and WKY in both schedules; this difference increased in FT 90-s. All strains started their drinking episodes later in FT 90-s than in FT 30-s. SHR produced longer episodes in FT 90-s relative to FT 30-s, and relative to other strains. Between strain differences: \* $p < 0.05$ , \*\* $p < 0.01$ . Within strain differences: # $p < 0.05$ , ## $p < 0.01$ .

respectively). Post hoc comparisons also revealed that SHRs produced longer episode durations in FT 90-s than in FT 30-s ( $p < 0.01$ ).

In short, Fig. 2 shows that, compared to control strains, SHRs produced more and longer drinking episodes that were more resilient to longer inter-food intervals. Despite this resiliency, SHR performance was sensitive to changes in inter-food interval: drinking episodes in SHR started later and lasted longer with longer inter-food intervals.

### 3.2. Effects on BERM parameters

Table 1 shows the goodness of fit of BERM. PVAFs were, in all cases, satisfactorily high, although the PVAF for WKY rats in FT 90-s was substantially lower than in any other strain  $\times$  schedule cell.  $\Delta$ AICc offered similar information. Nonetheless, even the worse fit of BERM provided a substantially superior fit than a single exponential distribution: BERM estimates were  $e^{\Delta\text{AICc}/2} = e^{2118}$  more likely than those from a single shifted exponential, given WKY performance in FT 90-s.

Fig. 3 shows fits of BERM to individual rat data. BERM fit the data adequately, displaying the usual broken-stick survival

**Table 1**  
Mean ( $\pm$ SEM) goodness-of-fit of BERM parameters.

	PVAF	$\Delta$ AICc <sup>a</sup>
FT 30-s		
SHR	0.84 (0.03)	8997 (1308)
WISTAR	0.76 (0.08)	5644 (1709)
WKY	0.69 (0.09)	5988 (1364)
FT 90-s		
SHR	0.93 (0.01)	17,247 (2725)
WISTAR	0.84 (0.05)	5095 (1983)
WKY	0.63 (0.15)	4237 (1584)

<sup>a</sup>  $\Delta$ AICc is the difference in likelihood between Equation 1 and a single-exponential model, after correcting for number of free parameters.  $\Delta$ AICc was computed for each rat and averaged across rats within each strain and schedule. Higher  $\Delta$ AICc is indicative of a higher likelihood of Eq. (1), given the data of each strain.

function of ILIs also observed in free-operant inter-response times (IRTs [24,25]). This suggests that when rats were involved in SIP episodes, ILIs were sampled from two exponential distributions: more than 80% of ILIs were sampled from a distribution with a short mean ( $1/w$  estimates in median rats ranged between 12 and 16 ms), and the remainder were sampled from a distribution with a long mean ( $1/b$  estimates in median rats ranged between 81 and 370 ms).

Fig. 4 shows mean ( $\pm$ SEM) BERM parameters ( $p$ ,  $w$ , and  $b$ ) and the mean number of bouts per episode, estimated from drinking episodes. Although mean  $p$  estimates were greater than .90 (which translates into 10 licks/bout) for all strains and schedules, there were significant differences in estimates among strains revealed by a significant strain effect [ $F(2, 32) = 20.796$ ,  $p < 0.01$ ]. SHRs generally produced lower  $p$  estimates than both WKY and Wistar ( $p < 0.01$  and  $p < 0.02$  respectively).

No significant differences in estimates of  $w$  were observed among strains or between schedules. During a drinking bout, all rats licked, on average, 3–6 times per second.


An analysis of estimates of  $b$  revealed a significant strain [ $F(2, 32) = 3.734$ ,  $p < 0.04$ ] and schedule effect [ $F(1, 32) = 5.063$ ,  $p < 0.03$ ]. SHRs produced licking bouts at a significantly higher rate than WKY rats ( $p < 0.04$ ), and rats generally produced licking bouts at a higher rate in FT 30-s than in FT 90-s. An analysis of estimates of the number of bouts per episode revealed a significant strain effect [ $F(2, 32) = 36.814$ ,  $p < 0.01$ ]. Post hoc comparisons showed that SHRs produced more bouts per episode than Wistar rats ( $p < 0.01$ ), and Wistars produced more than WKY rats ( $p < 0.01$ ).

In short, Fig. 4 shows that, compared to control strains, the drinking episodes of SHRs included more but shorter licking bouts. Within each bout, however, the rate of licking was very similar across strains.

## 4. Discussion

This study investigated the development and maintenance of SIP in SHR, Wistar and WKY rats in two FT schedules of food delivery. Results suggest a retardation in SIP acquisition in Wistar and WKY rats with respect to SHRs; the slowest acquisition was observed in WKY rats (Fig. 1). After 40 experimental sessions, when SIP performance stabilized, no significant differences in overall licking rate were observed among strains in FT 30-s. In FT 90-s, control strains hardly developed SIP. In contrast, SHRs maintained their adjunctive drinking despite a reduction in feeding rate. These results are consistent with previous reports [22,23].

The finding of differences in acquisition of SIP by the three strains of rats is interesting from a theoretical standpoint on adjunctive behaviour. WKY has been proposed as a model of anxiety vulnerability, post traumatic-stress and depression [32,33], and




**Fig. 3.** Sample fits of BERM (Eq. (1); continuous curves) to individual log-survival plots (the natural log of the probability of an LI greater than X) from selected SHR (top panels), Wistar (middle panels), and WKY rats (bottom panels). For each strain  $\times$  schedule (FT 30-s on the left, FT 90-s on the right) cell, the rat that produced the median lick/min was selected. Each panel includes BERM parameter estimates. BERM provided an adequate account of individual SIP performance.

there are theories of SIP that rely on its role as a coping strategy to mitigate the aversive conditions imposed by the intermittent presentation of food (e.g., Brett and Levine [34] observed reduced corticosterone levels as SIP training progressed). If SIP functions as an anti-anxiety behaviour, more licking should be expected in WKY than in the rest of rats. Results reported here were the opposite: WKY rats showed retardation in the acquisition of SIP in comparison to both SHR and Wistar rats, reaching with prolonged training (when developed) similar final levels of drinking. This learning deficit relates to other deficits observed in WKY rats under a variety of different behavioural tasks (e.g., [35]). Conversely, SHRs developed SIP quicker than Wistar controls, and this could reflect less competition for expression from other behaviours within inter-food intervals [22] and/or an increased hyperactivity [23]. Overall

the present results seem to reflect that the functionality of SIP is more in line with adjustments of behaviour to reinforcement intermittency (see [8,36]) rather than with an anxiety-reduction function.

After 40 training sessions, not all rats acquired SIP above prandial levels, and were therefore excluded from further analysis. All excluded rats were Wistar and WKY; none were SHR. Even after excluding low responders, however, selected Wistar and WKY rats produced drinking episodes at a much lower rate than SHRs (Fig. 2A). Furthermore, the episode durations of SHRs were longer than those of controls in FT 90-s (Fig. 2C). Taken together, these results suggest that licking behaviour was stronger in SHR than in control strains, particularly in FT 90-s. Despite this difference, all selected rats produced enough ILLs in both schedules to



**Fig. 4.** Mean ( $\pm$ SEM) BERM parameters (Eq. (1)) estimated from drinking episodes. The figure A shows estimates of  $p$ , the probability of continuing a bout after every lick. The figure B shows estimates of  $w$ , the within-bout licking rate. The figure C shows estimates of  $b$ , the bout-initiation rate. The figure D shows estimations of the average number of bouts per episode, computed as  $w(bD+1)/(w+Lb)$ , where  $D$  is the mean episode duration, and  $L$  is the mean bout length  $= 1/(1-p)$ . SHR produced shorter episodes (lower  $p$ ) than WKY and Wistar rats. SHR produced bouts more frequently than WKY. SHR produced more bouts per episode than Wistar rats, and Wistar rats more than WKY in both schedules. Between strain differences: \* $p < 0.05$ , \*\* $p < 0.01$ . Within strain differences: # $p < 0.05$ , ### $p < 0.01$ . - $p > 0.05$ .

demonstrate a superior fit of BERM relative to a more parsimonious alternative model (Table 1 and Fig. 3).

The adequate fit of BERM suggests that ILIs are sampled from two exponentially distributed populations of ILIs, one with a shorter mean that corresponds to within-bout ILIs, and one with a longer mean that separates these bouts. This pattern of behaviour has 3 parameters: the proportion of ILIs sampled from the short distribution ( $p$ ); which covaries positively with mean bout length), the within-bout licking rate ( $w$ ), and the bout-initiation rate ( $b$ ).

The pattern of differences in parameter estimates across strains is remarkably similar to the pattern of differences drawn from IRTs in variable-interval (VI) schedules of food reinforcement [4].

This similarity is particularly striking when ILI and IRT distribution parameters are drawn from adult rats [4]. Under these conditions, WKY produce very long bouts (i.e., high  $p$ ) of both operant and adjunctive behaviour, compared to other strains (Fig. 4A). Both operant and adjunctive bout lengths do not appear to be significantly affected by rate of reinforcement. No clear differences in within-bout response rate are observed across strains or schedules in either operant or adjunctive behaviour (Fig. 4B). SHR produces substantially more operant and adjunctive bouts than Wistar, and Wistar produces more than WKY (Fig. 4C). Taken together, these data suggest that differences across strains in the patterns of food-elicited behaviour are not dependent on response-food contingencies, even though bout-initiation rates are sensitive to the frequency of food delivery. The similarity of results across studies is consistent with the notion that the laws that govern adjunctive behaviours, such as SIP, are the same that govern operant behaviour [36].

Hill et al. [4] infer from operant performance that *operant hyperactivity* (the inordinately high rate of responding at low rates of reinforcement) of SHR is due to a combination of steep reinforcement gradient (supporting shorter bouts) and, possibly, an enhanced responsiveness to reinforcement (supporting more frequent bouts). The present results extend these inferences to situations in which incentives are not contingent on behaviour. Thus, operant hyperactivity in SHR may be thought of as belonging to a wider class of hyperactive behaviour expressed by SHR, which may be called *incentive-elicited hyperactivity*.

## References

- [1] Taylor D, Lincoln AJ, Foster SL. Impaired behavior regulation under conditions of concurrent variable schedules of reinforcement in children with ADHD? *J Atten Disord* 2010;13(4):358–68.
- [2] Sagvolden T. Behavioral validation of the spontaneously hypertensive rat (SHR) as an animal model of attention-deficit/hyperactivity disorder (AD/HD)? *Neurosci Biobehav Rev* 2000;24(1):31–9.
- [3] Sanabria F, Killeen PR. Evidence of impulsivity in the Spontaneously Hypertensive Rat drawn from complementary response-withholding tasks. *Behav Brain Funct* 2008;4:11L 17.
- [4] Hill C, Herbst K, Sanabria F. Characterizing operant hyperactivity in the Spontaneously Hypertensive Rat. *Behav Brain Funct* 2012;8(5):1–15.
- [5] Falk JL. Production of polydipsia in normal rats by an intermittent food schedule. *Science* 1961;133:195–6.
- [6] Falk JL. Schedule-induced polydipsia as a function of fixed interval length. *J Exp Anal Behav* 1966;9:37–9.
- [7] Falk JL. Control of schedule-induced polydipsia: type, size, and spacing of meals. *J Exp Anal Behav* 1967;10(2):199–206.
- [8] Falk JL. The nature and determinants of adjunctive behavior. *Physiol Behav* 1971;6:577L 88.
- [9] Staddon JER. Schedule-induced behavior. In: Honig WK, Staddon JER, editors. *Handbook of operant behavior*. Englewood Cliffs, NJ: Prentice-Hall; 1977. p. 125–52.
- [10] Riley AL, Wetherington CL. The differential effects of naloxone hydrochloride on the acquisition and maintenance of schedule-induced polydipsia. *Pharmacol Biochem Behav* 1987;26(4):677–81.
- [11] Gilpin NW, Badia-Elder NE, Elder RL, Stewart RB. Schedule-induced polydipsia in lines of rats selectively bred for high and low ethanol preference? *Behav Genet* 2008;38(5):515–24.
- [12] Cardona D, López-Grancha M, López-Crespo G, Nieto-Escamez F, Sánchez-Santed F, Flores P. Vulnerability of long-term neurotoxicity of chlorpyrifos: effect on schedule-induced polydipsia and a delay discounting task. *Psychopharmacology (Berl)* 2006;189:47L 57.
- [13] López-Grancha M, López-Crespo G, Sánchez-Amate MC, Flores P. Individual differences in schedule-induced polydipsia and the role of gabaergic and dopaminergic systems. *Psychopharmacology (Berl)* 2008;197:487L 498.
- [14] Myracle A, Lopez-Grancha M, Flores P, Glowa J, Riley AL. Differential effects of morphine and LiCl on schedule-induced polydipsia? *Pharmacol Biochem Behav* 2005;80(1):195–202.
- [15] Plat B, Beber C, Schechter L, Rosenzweig-Lipson S. Schedule-induced polydipsia: a rat model of obsessive-compulsive disorder. *Curr Protoc Neurosci* 2008;9:9–27.
- [16] Van Kuyck K, Brak K, Das J, Rizopoulos D, Nuttin B. Comparative study of the effects of electrical stimulation in the nucleus accumbens, the mediodorsal thalamic nucleus and the bed nucleus of the stria terminalis in rats with schedule-induced polydipsia. *Brain Res* 2008;1201:93L 99.

- [17] Moreno M, Flores P. Schedule-induced polydipsia as a model of compulsive behavior: neuropharmacological and neuroendocrine bases. *Psychopharmacology (Berl)* 2012;219:647–59.
- [18] Moreno M, Gutierrez-Ferre VE, Ruedas L, Campa L, Sunol C, Flores P. Poor inhibitory control and neurochemical differences in high compulsive drinker rats selected by schedule-induced polydipsia? *Psychopharmacology (Berl)* 2012;219(2):661–72.
- [19] Flores P, Sánchez-Kuhn A, Merchán A, Vilches O, García-Martín S, Moreno M. Schedule-induced polydipsia: searching for the endophenotype of compulsive behavior. *World J Neurosci* 2014;4:253L 260.
- [20] Hawken ER, Beninger RJ. The amphetamine sensitization model of schizophrenia symptoms and its effect on schedule-induced polydipsia in the rat? *Psychopharmacology (Berl)* 2013;231(9):2001–8.
- [21] Hawken ER, Delva NJ, Beninger RJ. Increased drinking following social isolation rearing: implications for polydipsia associated with schizophrenia. *PLOS ONE* 2013;8(2):e56105.
- [22] Íbias J, Pellón R. Schedule-induced polydipsia in the Spontaneously Hypertensive Rat and its relation to impulsive behaviour. *Behav Brain Res* 2011;223:58L 69.
- [23] Íbias J, Pellón R. Different relations between schedule-induced polydipsia and impulsive behaviour in the Spontaneously Hypertensive Rat and in high impulsive Wistar rats: questioning the role of impulsivity in adjunctive behaviour. *Behav Brain Res* 2014;271:184L 194.
- [24] Shull RL, Grimes JA, Bennett JA. Bouts of responding: the relation between bout rate and the rate of variable-interval reinforcements. *J Exp Anal Behav* 2004;81:65L 83.
- [25] Brackney RJ, Cheung TH, Neisewander JL, Sanabria F. The isolation of motivational, motoric, and schedule effects on operant performance: a modelling approach. *J Exp Anal Behav* 2011;96:17L 38.
- [26] Brackney RJ, Cheung TH, Herbst K, Hill JC, Sanabria F. Extinction learning deficit in a rodent model of attention-deficit hyperactivity disorder. *Behav Brain Funct* 2012;13:8L 59.
- [27] Kessel R, Lucke RL. An analytic form for the interresponse time analysis of Shull, Gaynor, and Grimes with applications and extensions. *J Exp Anal Behav* 2008;90:363L 386.
- [28] Shull RL, Gaynor ST, Grimes JA. Response rate viewed as engagement bouts: effects of relative reinforcement and schedule type? *J Exp Anal Behav* 2001;75(3):247–74.
- [29] Fleishman AI. Confidence intervals for correlation ratios. *Educ Psychol Meas* 1980;40:659–70.
- [30] Burnham KP, Anderson DR. Model selection and multimodel inference: a practical information-theoretic approach. 2nd ed. New York, NY: Springer-Verlag; 2002.
- [31] Cheung TH, Neisewander JL, Sanabria F. Extinction under a behavioral microscope: isolating the sources of decline in operant response rate? *Behav Process* 2012;90(1):111–23.
- [32] Solberg L, Losee S, Turek F, Redei E. Altered hormone levels and circadian rhythm of activity in the WKY rat, a putative animal model of depression. *Am J Physiol Regul Integr Comp Physiol* 2001;281:786L 94.
- [33] Will CC, Aird F, Redei EE. Selectively bred Wistar-Kyoto rats: an animal model of depression and hyper-responsiveness to antidepressants. *Mol Psychiatry* 2003;8:925–32.
- [34] Brett LP, Levine S. The pituitary–adrenal response to minimized schedule-induced drinking. *Physiol Behav* 1981;26:153L 158.
- [35] McAuley JD, Stewart AL, Webber ES, Cromwell HC, Servatius RJ, Pang KCH. Wistar-Kyoto rats as an animal model of anxiety vulnerability: support for a hypervigilance hypothesis. *Behav Brain Res* 2009;204:162–8.
- [36] Killen P, Pellón R. Adjunctive behaviours are operants. *Learn Behav* 2013;41:1L 24.